

Article ID: 1006-8775(2019) 01-0102-12

URBAN HEAT ISLAND EFFECT AND ITS CONTRIBUTION TO OBSERVED TEMPERATURE INCREASE AT WUHAN STATION, CENTRAL CHINA

JIA Wen-qian (贾文茜)¹, REN Guo-yu (任国玉)^{1,2}, SUONAN Kan-zhuo (索南看卓)¹, ZHANG Pan-feng (张盼峰)¹,
WEN Kang-min (温康民)¹, REN Yong-jian (任永健)³

(1. Department of Atmospheric Sciences, China University of Geosciences, Wuhan 430070 China; 2. Laboratory for Climate Studies, China Meteorological Administration, National Climate Center, Beijing 100081 China;
3. Wuhan Regional Climate Center, Wuhan 430074 China)

Abstract: Based on an in-homogeneity adjusted dataset of the monthly mean temperature, minimum and maximum temperature, this paper analyzes the temporal characteristics of Urban Heat Island (UHI) intensity at Wuhan Station, and its impact on the long-term trend of surface air temperature change recorded during 1961–2015 by using an urban-rural method. Results show that UHI effect is obvious near Wuhan Station in the past 55 years, especially for minimum temperature. The strongest UHI intensity occurs in summer and the weakest in winter. For the period 1961–2004, UHI intensity undergoes a significant increase near the urban station, with the increase especially large for the period 1988–2004, but the last 10 years witness a significant decrease, with the decrease in minimum temperature being more significant than that of maximum temperature. The annual mean urban warming and its contribution to overall warming are 0.18 C/10yr and 48.8% respectively for the period 1961–2015, with a more significant and larger urbanization effect seen in T_{\min} than T_{\max} . Thus, a large proportion warming, about half of the overall increase in annual mean temperature, as observed at the urban station, can be attributed to the rapid urbanization in the past half a century.

Key words: data in-homogeneity; UHI; urban warming; climate change

CLC number: P467 **Document code:** A

doi: 10.16555/j.1006-8775.2019.01.010

1 INTRODUCTION

Climate change research is mainly based on continuous and high-quality observational records. Due to the relocation of observation sites, replacement of observation instruments and changes in the time system, however, inhomogeneities have occurred in the observational data series to varying degrees, and they can cause a significant bias in the analysis of surface climate variables trends for individual stations or a small area. Furthermore, the gradually increasing Urban Heat Island (UHI) effect near the observational sites becomes the biggest systematic bias for the long-term climate change monitoring and detection in that it directly causes the trend of surface air temperature (SAT) records to be overestimated in many developing regions of the world.

At present, there exists a major divergence of views in the international climatological community on the impact degree of urbanization effect on the SAT series. It is generally held that UHI effect is small, and it has not

surpassed 0.05°C in the past 100 years on a global average, much lower than the optimal estimation of the global average annual mean SAT change of 0.8 °C (Jones et al.^[1]; Peterson^[2]; Parker^[3]; Lawrimore et al.^[4]). Nevertheless, some researches for several regions have shown that the UHI effect may play a significant role in the regional SAT trend, which should be paid more consideration to (Karl et al.^[5]; Hansen et al.^[6]; Kalnay and Cai^[7]; Zhou et al.^[8]; Zhou and Ren^[9]; Ren et al.^[10]).

The studies of the UHI effect on SAT records have been conducted for big cities and large regions in recent years, and stimulating conclusions have been drawn thereby. For example, the investigation of UHI effect on Beijing and Shijiazhuang stations in northern China has shown that the UHI Intensity(UHII) near the observational sites is stronger at night and in winter than during the daytime and in summer (Xie et al.^[11]; Zheng et al.^[12]; Ren et al.^[13]; Yang et al.^[14]; Bian et al.^[15]); Deng et al. pointed out that the UHII at night in Shanghai was the strongest in autumn/winter, while the UHII during daytime was relatively stronger for the spring and plum rain season of early summer, and lowest in autumn^[16]. Meanwhile, studies also found that change trends of annual mean SAT observed at meteorological stations were clearly influenced by urban warming. For example, Li et al. showed that the annual mean SAT trend brought about by urbanization for southern coastal stations in China is higher than the background trend^[17]; Zhou and Ren found that the contribution of urban warming to the overall warming in the northern

Received 2018-01-31; **Revised** 2018-11-13; **Accepted** 2019-02-15

Foundation item: National Natural Science Foundation of China (41575003), National Key R&D Program of China (2018YFA0605600)

Biography: JIA Wen-qian, M.S., primarily undertaking research on urban climate.

Corresponding author: REN Guo-yu, e-mail: guoyoo@cma.gov.cn

China region was 37.9% during 1961–2000^[9], and it even reached 71% at Beijing station over the same period (Chu and Ren^[18]); Urbanization-induced SAT trend reached 0.19°C/10yr during 1962–2009 for Shijiazhuang station, accounting for 68% of all warming trend^[15].

Wuhan, the capital city of Hubei Province, has a built-up area of approximately 400 km² and a population of over 5 million, experiencing a rapid development and urbanization process. Wuhan city locates in the Jiangnan Plain (Fig.1) and always experiences severe heat wave and rainstorms in summer, which probably has been aggravated by urbanization effect during the past two decades. UHI effect has been remarkable at Wuhan station since the 1980s, especially in winter, and the minimum temperature witnesses the highest contribution of urban warming to overall warming (Ren et al.^[13]; Wu et al.^[19]); Chen et al. studied the UHI change trend for Wuhan station from 1960 to 2005, and found that the annual mean urban warming and contribution rate reached 0.235°C/10yr and 60.4% respectively^[20]. However, the previous researches only rely on simple criteria to select reference stations, and the selected rural stations may have been gradually affected by UHI effect with the development of small cities and towns themselves. There is a need to adopt a more objective method to select reference stations, and to assess whether the analysis results obtained by the new method are consistent with previous ones. In addition, it has been found that the temperature and sunshine hours in Wuhan decreased in recent 10 years (Cao^[21]; Fyfe^[22]), which is consistent with the global warming slowdown, and the relative humidity (RH) experienced a rise within a certain degree after 2007 (Suonan et al.^[23]). It is not clear whether UHI effect for Wuhan station will have some new features when the data is updated to present to include the period of climate warming slowdown.

In this paper, in-homogeneity adjustment is made for the SAT data updated to 2015. We analyze the climatological characteristics of UHI effect and the change of UHI with time near Wuhan station by using the homogeneous data from the urban station and two rural stations. Meanwhile, we estimate the urbanization effect on the long-term SAT change trend of the Wuhan station. The results of this paper are helpful for a further understanding of temporal characteristics of UHI effect, as well as nature and extent of urbanization-induced bias of SAT observation records for Wuhan station.

2 DATA AND METHODS

The monthly mean, minimum and maximum SAT data from 13 stations in Wuhan for the time period 1961–2015 and the MOD11A2 8-day LST product at 1-km spatial resolution in 0.02K unit from July 28 to August 4 in 2016 are used. The SAT are obtained from the National Meteorological Information Center of the China Meteorological Administration, which have already been quality controlled and the inhomogeneities caused by the shift from artificial observation to automatic observation

have been corrected. The SAT data at Caidian station missed in June, July and August in 1968 was replaced by monthly mean SAT data for 55 years.

Gallo et al.^[24] used the vegetation index obtained from NOAA/AVHRR to explain the temperature difference between urban and rural areas. Ren and Ren used MODIS data to select 113 SAT reference stations from 672 national reference climate stations and national basic meteorological stations of mainland China, pointing out that the method could be adopted for the adjustment study on the urbanization bias of the currently used air temperature records of surface climate stations^[25]. Compared to the other approaches used for classifying climatic stations, the remote sensing method does not rely so much on social and economic data. It is therefore feasible to use MODIS products to determine the regional boundaries of UHI-affected areas (Zhang^[26]).

The remote sensing image quality is affected by cloud cover. Hubei is controlled by the Subtropical High in late July and the early August, and the weather condition is better. Therefore, the average LST from July 28 to August 4 in 2016 is used to determine the spatial distribution of ground surface temperature around the stations. The isothermal pattern of LST could reflect the temperature gradient from urban centers to the suburbs that are less influenced by the UHI effect^[25]. Winkler et al. proposed that the boundary of the area affected by the UHI effect can be represented by the outermost closed isotherm of the annular temperature field in a city or town^[27]. In this paper, the outermost isotherm circle is 33.85 °C and it can be taken as the boundary of the UHI-affected area in the Wuhan city (Fig.1).

The urban stations and the reference (rural) stations are selected according to the specific location of the station in the brightness temperature field. The stations which are located within the isotherm 33.85°C are regarded as urban stations. The selection of the reference stations follows the following principles: (1) outside the closed contour 33.85 °C of surface temperature; (2) the distance from the boundary of built-up areas within 30km to ensure the same large-scale circulation and climate background with the urban stations; (3) elevation difference with the urban stations less than 20m, avoiding the influence brought by height difference; (4) the micro-environment around the rural stations is open.

The selection of rural stations is particularly important for studying the urban heat island effect. We therefore chose a more objective and strict method for the reference stations selection. Compared with other approaches to classify climatological stations, the remote sensing (RS)-based method is much less dependent on social or economic data, and RS data is also relatively continuous, with high spatial comparability and objectivity.

In addition to ensuring that all historical stations are outside the outermost contour when selecting a reference station, the distance to the city station, the altitude

difference, and the environment around the sites are also taken into consideration. The distance and elevation difference are limited to ensure in the same climate background with urban station, reducing the errors of results as far as possible. In addition, many natural landscapes, such as lakes, basins and hills, also affect the surface thermal structure, and may result in the closed isotherms around the stations. These have also been considered when determining the reference stations.

The locations of all historical observation sites for Wuhan station are located within the closed circle of 33.85°C (Fig.1), and there has been a weak but clear urban heat island effect at urban stations (Wang^[28]). We therefore regard Wuhan station as a representative urban

station; the rural stations of all historical observation outside the closed circle of 33.85°C are Huangpi, Hanchuan and Anlu stations. The Anlu station is too far from Wuhan station and the elevation difference from the urban station is more than 20m (Table 1), so it is not taken as a reference station, and the relocation process of Anlu station is not shown in Fig.1. From Google Earth images(Fig.2), we find that the locations of remaining rural stations are at the edge of the built-up areas, with little population and few buildings. These two stations have not yet been significantly affected by nearby settlement and the UHI effect, and they can therefore be regarded as rural stations.




Figure 1. The location of 13 stations and average LST isotherm distributions in Wuhan. Abbreviations of the stations' names are AL for Anlu, HA for Hong'an, YM for Yunneng, XG for Xiaogan, HP for Huangpi, XZ for Xinzhou, HC for Hanchuan, WH for Wuhan, CD for Caidian, JX for Jiangxia, HG for Huanggang, EZ for Ezhou, and JY for Jiayu. The black line in the graph is the boundary of Wuhan city, and the relative position of Wuhan is shown in the illustration. The three enlarged figures on the right represent the relocation process of selected urban and rural stations, and the arrows show the relocation direction.

Table 1. Information of the stations used in the study.

Station name (Code)	Altitude(m.a.s.l.)	Distance from Wuhan station (m)	The time and distance of Relocations since 1961(km)
Wuhan(57494)	23.6	0	1994.4(0.08); 2010.1(8)
Hanchuan(57486)	27.0	26110	1964.1(6); 1970.11(3)
Huangpi(57491)	31.7	39100	
Anlu(57388)	55.5	84110	




Figure 2. Station and the surrounding landscapes as seen from Google Earth picture: (a) Hanchuan (No.57486), and (b) Huangpi (No. 57491).

The breakpoints of data series caused by relocations are examined for annual mean (T_{mean}), minimum (T_{min}) and maximum (T_{max}) SAT using E-P method (Easterling and Peterson^[29-30]; Li et al.^[31]). The autocorrelation problem needs to be considered in the in-homogeneity test. Annual mean SAT data series is less affected by the seasonal variability, reducing data series autocorrelation to a large extent (Cao and Yan^[32]). Therefore, in-homogeneity test and correction are based on the annual temperature data series in this paper.

Locations of the breakpoints could be determined based on whether the values of t statistics exceed the significance level (Wei^[33]). An adjustment could be made if the breakpoints could be proved to be real with reference to the metadata. The difference between the five-year average before and after the breakpoints is used as the adjustment value, and if the data is less than 5 years

before and after the breakpoints, the longest time is taken to calculate the difference. Then the annually averaged adjustment values are linearly interpolated to each of the months. Here, $n=55$, the length of the sub-series was 3 years, and the given significance level $\alpha=0.05$.

After an investigation of historical data, it is found that the relocation of Wuhan station in 2010 with a move of about 8 km has a significant impact on temperature series. The T_{mean} and T_{min} series both witnessed big breakpoints in 2010 by using moving t -test method. The breakpoints therefore for the T_{mean} and T_{min} data series are adjusted in the study. To ensure data uniformity, the T_{max} series also has been corrected despite the breakpoint in 2010 is undetectable. In the process of adjustment, we correct the SAT data after station moving by adding the adjustment values to better keep the UHI effect recorded at the original site (Wang et al.^[34]) (Fig.3).




Figure 3. A comparison of the series of annual mean, minimum and maximum temperature before and after homogenization at Wuhan station.

Hanchuan station has been moved two times since 1961, but the data series has no detectable breakpoint probably because the relocation time was too early, and the data series has not been adjusted. Huangpi station has not been moved since its establishment, and this enables it to be an ideal reference station for examining the urbanization effect.

In the process of SAT data adjustment for Wuhan station, especially for the T_{mean} and the T_{min} , the corrected values are positive, indicating that T_{mean} and T_{min} originally drop sharply since 2010, which is largely due to the Wuhan station moving from the urban area to the suburbs at that time. There is a false cooling phenomenon in SAT series of Wuhan station caused by the relocation, and the warming trend of the corrected temperature series is more obvious than that of the original one (Fig.3). The homogenization has regained the UHI effect and urban warming trend in the SAT series (Zhang et al.^[35]).

UHII at Wuhan station is defined as the SAT difference between the urban and rural stations. The rural SAT or reference series is the average SAT of the two rural (reference) stations. Linear trends of the UHII series are obtained by using the least square method and the significance level of linear trend is examined by t -test method. The difference of SAT linear trends between urban and rural stations is equal to the linear trend of the UHII series, and they are defined as urban warming or urbanization effect. Contribution of urbanization effect or urbanization contribution is the proportion of the

statistically significant urbanization effect on the overall trend of SAT at the urban station^[10,13].

In practical calculations, the urbanization contribution may exceed 100%, probably brought by unknown local anthropogenic factors for rural stations in a few cases. In this case, it is adjusted to 100%. As indicated in the definition, if the urban warming is not statistically significant, the urbanization contribution will not be calculated or analyzed.

3 RESULTS AND DISCUSSION

3.1 Climatological characteristics of UHII

Annual and seasonal mean UHII at Wuhan station for two periods of 1961–2015 and 1988–2015 are examined. As described later, UHII near Wuhan station is relatively weak before 1988, and a significant increase occurs afterward (Fig.5).

The annual mean UHII at Wuhan station is 0.39°C for the period 1961–2015, the largest seasonal mean UHII is 0.47°C in summer, followed by 0.41°C in spring, 0.40°C in autumn, and 0.32°C in winter, but overall the seasonal difference is not significant (Table 2). The minimum temperature UHII is higher than the maximum temperature UHII, indicating that the UHII in Wuhan is stronger at night than it is in the daytime. UHII has an obvious diurnal non-symmetry, which is consistent with most previous research results of the megacities in China (Xie et al.^[11]; Zheng et al.^[12]; Bian et al.^[15]; Zhou and Zhang^[36]).

Table 2. Annual and seasonal mean, minimum and maximum temperature UHII for time periods 1961–2015 and 1988–2015 at Wuhan station (°C).

Period	Temperature	Spring	Summer	Autumn	Winter	Year
1961–2015	Mean	0.41	0.47	0.40	0.32	0.39
	Minimum	0.56	0.61	0.63	0.40	0.54
	Maximum	0.35	0.41	0.21	0.29	0.32
1988–2015	Mean	0.67	0.81	0.73	0.58	0.69
	Minimum	0.99	1.14	1.26	0.95	1.07
	Maximum	0.44	0.51	0.29	0.33	0.39

In the past 28 years, the seasonal feature of the mean UHII for Wuhan station is consistent with that of 1961–2015, the largest in summer and smallest in winter. The difference is that the three kinds of temperature UHII is larger during the recent period, with the minimum temperature UHII being the most obvious, almost two times that during the 1961–2015 period, followed by the mean temperature UHII, and the maximum temperature UHII change is the smallest. It shows that the UHII near Wuhan station has been enhanced since the late 1980s, which well corresponds to the rapid urbanization process in the city. The population and built-up areas in Wuhan had a rapid increase after the late 1980s and early 1990s, leading to an accelerating urbanization process (Xiao et

al.^[37]).

Both throughout the entire period and in the recent 28 years, the seasonal mean temperature and maximum temperature UHII are the largest in summer, and the minimum temperature UHII is the strongest in fall. These features are different from the UHI effect of the metropolis in northern China in which the strongest UHII occurs in winter and the weakest in summer or spring (Xie et al.^[11]; Zheng et al.^[12]; Yang et al.^[14]; Bian et al.^[15]; Bai et al.^[38]). This may be mainly due to the smaller solar elevation angle in winter (the zenith angle is larger), and the frequent temperature inversion and stable weather phenomenon controlled by the dry and cold Siberia air mass in winter in the north. Meanwhile, the humidity in

lower atmosphere and cloudiness in northern China is usually less and anthropogenic heat release due to the heating is large in winter (Chen and Shi^[39]). In contrast to winter, summertime in the north has a larger solar elevation angle (smaller zenith angle), and is more frequently affected by the southwest and southeast air currents, with more cloudiness and precipitation, and cooling energy consumption and artificial heat release in urban areas is smaller, leading to a generally lower UHI effect; the average wind speed in spring in the north is usually stronger, which is also against to the UHI formation and development.

Wuhan features an inland subtropical monsoon humid climate, with hundreds of lakes scattering in urban areas. In winter, the relative humidity and water vapor content of lower atmosphere becomes larger, and the anthropogenic heat release from urban heating is less, which may be the main reasons for the lower UHII in the cold season. In summer, the atmospheric humidity is high, with more cloudiness and precipitation, and the solar elevation angle is large (the zenith angle is small). These will not benefit the formation and development of UHI. However, the average wind speed in summer is weaker, and stable atmospheric layer and more sunny days prevail, because the city is controlled by the subtropical high in midsummer season^[28]. Besides, the use of air conditioning in the summer is becoming more common every year, with large anthropogenic heat release into the urban boundary layer. All of these will benefit the enhancement of the UHII in the urban areas during

summertime. The weather in autumn is sunny with little cloud and the wind speed is small. In addition, the heat storage by abundant urban lakes makes boundary layer atmosphere warmer, enhancing the UHII. A higher seasonal mean UHII in autumn therefore results.

The seasonal variation feature of UHII at Wuhan station is like the other southern cities, such as Shanghai city and Guangzhou city, with larger UHII in autumn for mean and minimum temperature and significantly more obvious UHI phenomenon in spring and summer for maximum temperature. There is a big difference for the UHII of summer and winter for mean and minimum temperature between Wuhan station and other southern cities, however, and this may be related to the different underlying surface, geographical position and other factors (Deng et al.^[16]; Zhou and Zhang^[36]; Xu^[40]).

The seasonal variations of monthly mean UHII in the past 55 years and the past 28 years are similar (Fig.4). The mean temperature UHII and minimum temperature UHII are the lowest in January, and then they gradually increase. The monthly mean temperature UHII peaks in July, and then slowly weakens, while the peak of minimum temperature UHII appears in September, and higher in November for the recent 28 years. The monthly mean maximum temperature UHII has a different seasonal cycle, with the strongest in July and the weakest in October. It is thus obvious that the difference between the maximum and minimum temperature UHII in autumn is the largest in a year.




Figure 4. Seasonal variation of monthly mean, minimum and maximum temperature UHII for time periods 1961-2015 and 1988-2015 at Wuhan station (°C).

The fact that maximum temperature UHII shows a downward trend after July can be explained by the increase in rainfall and cloudiness significantly accompanying the retreat of the Subtropical High, with the suburbs and the city receiving similarly small amount of solar radiation during the daytime. The reduced anthropogenic heat release after mid-August might be another reason. In addition, large areas of lake and wetland in urban make night warmer and day cooler, with the latter offsetting the maximum temperature UHI effect to certain extent.

3.2 Urban warming and its contribution to overall

warming

Figure 5 provides the long-term change in annual mean, minimum and maximum temperature UHII at Wuhan station. Annual mean and minimum temperature UHII display a consistent inter-annual and decadal variation, showing that UHII gradually increase since the early 1960s, especially for minimum temperature UHII as high as 33%, reaching peak at around 2005 and dropping afterward. The maximum temperature UHII experiences a certain degree of decline in the 1960s, remains stable in the 1970s and 1980s, and increases slowly from the 1990s to 2004.




Figure 5. Annual mean, minimum and maximum temperature UHII (polygonal lines) and their trends (straight lines) at Wuhan station.

The UHII variations in the three kinds of temperatures are all characterized by lower values from the 1960s to the late 1980s, higher values after 1988. In general, the three kinds of temperatures UHII show an upward trend in the past 55 years, with an obvious rising in 1961–2004, the fastest increase ($0.53^{\circ}\text{C}/10\text{yr}$ for annual mean minimum temperature) in 1988–2004, and a decrease in 2005–2015. Therefore, the long-term trends of UHII can be examined for four different time periods of 1961–2015, 1961–2004, 1988–2004 and 2005–2015.

Figure 6 shows the urbanization-induced change in annual mean, minimum and maximum SAT, and the urbanization contribution to the overall warming at Wuhan station for the four periods. For the period 1961–2015, the trend of annual mean UHII and the urbanization contribution are $0.18^{\circ}\text{C}/10\text{yr}$ and 48.8% respectively. A more evident urban warming and contribution occur in T_{\min} than T_{\max} . For the seasons, urban warming and contribution for T_{mean} are both highest in summer, followed by autumn. The largest urban warming of $0.37^{\circ}\text{C}/10\text{yr}$ occurs in autumn T_{\min} . Seasonal mean warming observed in summer T_{mean} can be entirely accounted for by the enhanced UHI effect.

Seasonal mean urban warming and urbanization contribution for 1961–2004 are like those for the period 1961–2015, showing more evident change in summer and autumn. The difference is that contribution of urban warming in summer T_{\max} is a negative value of -77.9% , implying that the background maximum SAT change during this period is characterized by a cooling trend and the urban warming is making an opposite contribution to the decrease at Wuhan station.

The urban warming during 1988–2004 is larger and more significant in comparison to it is in the periods 1961–2004 and 1961–2015. Therefore, more significant urbanization effect at Wuhan station occurs during the late 20th century when urbanization and economic growth of China are unprecedented in history. The annual and seasonal mean, minimum and maximum temperature UHII trends are significantly greater than the other several stages, with more evident urban warming for summer and autumn T_{mean} . However, the urbanization contribution for summer T_{mean} becomes weaker relative to the periods 1961–2015 and 1961–2004. Autumn T_{\min} witnesses the most significant urban warming reaching as high as $1.00^{\circ}\text{C}/10\text{yr}$, and the urbanization contribution as high as 65%, showing the warming observed at autumn night can largely be attributed to the urbanization effect.

Seasonal mean maximum temperature UHII trends and urbanization contribution during 1988–2004 are much larger than those in any other periods. This is especially true for summer, with the urbanization contribution even reaching 100%, indicating an entire contribution from the increased UHII to the positive trend of summer mean maximum temperature at the urban station (Fig.6).

It is worth noting that a relatively small urban warming but quite significant urbanization contribution at Wuhan station registers in winter, which is completely contrary to the results of the megacities in northern China^[10,15,18], where the UHII trends are the largest in winter, and the urbanization contribution to the overall seasonal warming is often smaller. This may be related to the lower latitude, wetter and hotter climatic condition




Figure 6. Urbanization effects (solid) and the urbanization contribution (blank) of seasonal and annual mean, minimum and maximum surface air temperature at Wuhan station.

and larger proportion of wetland area in Wuhan city compared to those northern megacities. The anthropogenic heat-release difference of winter heating and summer cooling between northern and southern cities, and the different background circulation fields and multi-decadal climate variability, would be also significant to the contrary seasonality of the urbanization effects.

It is worth noting that the rural stations unaffected by urbanization effect can hardly be found in central China. The rural stations used in this paper are currently available stations that are less affected by urbanization. Therefore, there is a certain degree of urbanization impact in the rural station temperature series. As a result, urban warming and their contributions to the overall warming in different time periods as given in this work for Wuhan station could be regarded as the lowest estimates.

China has witnessed a rapid urbanization process in the past half century, especially the past 30 years. From 1978 to 2015, the urbanization rate of the country increases from 17.9% to 50.0%. This phase is also accompanied by the obvious increase in urban heat island effect around the climatic stations^[10]. Through calculation, the urban land and construction land increase by 58%

from 1988 to 2004 in Wuhan area, with the most significant urban warming of Wuhan station as high as 0.53°C /10yr, indicating that the rapid expansion of the city is the main reason for the increase of UHI effect recorded in the SAT series.

Unlike T_{mean} and T_{max} , T_{min} at Wuhan station undergoes a significantly higher urban warming in the above three time periods (1961–2015, 1961–2004 and 1988–2004). This indicates that the UHI effect is more obvious at night than the daytime, which is mainly due to the greater thermal admittance in the urban area than that of the suburbs, leading to a slower temperature drop in urban after sunset. The urban buildings, with many urban street canyons, generally absorb more solar radiation than the suburbs during the daytime. There are downward long-wave radiations of walls and eaves in addition to atmospheric counter-radiation, coupled with the relatively low wind speed, and the heat is not easily dissipated at night in the urban area. All of these make the nighttime UHI effect higher than daytime, and the UHI effect at night will become more obvious with the development of the city, resulting a more rapid increase in T_{min} than T_{mean} and T_{max} .

During the period from 2005 to 2015, the annual and

seasonal mean, minimum and maximum temperature UHII trends are negative, indicating that the UHII for Wuhan station is weakening in the past 10 years. The descending rates of urban warming for annual and seasonal mean T_{\min} are much higher than the T_{\max} except for the summer. Although the UHII is weakening in the last 10 years, the urban contribution is positive and still large. This may be related to the fact that the background SAT is experiencing a decrease during the period of climate warming slowdown, and the urban station undergoes a more rapid decrease than the rural stations do (Fig.7). The relatively slower decrease in SAT at the rural stations may be caused by the increase of built-up areas around the observational sites in recent 10 years. The rural stations may therefore have been affected by urbanization. It is highly possible that the rural stations have become unrepresentative after 2005 due to the urbanization effect in these originally suburban areas.

The annual mean T_{mean} , T_{\min} and T_{\max} in urban and rural stations have decreased since 2005 (Fig. 7), which is consistent with the trend of temperature changes in China. Xu pointed out that the annual mean temperature shows a certain degree of downward trend from 2005–2012^[40]. The warming hiatus since about 2000 is especially obvious in north China (Wang et al.^[41]). The temperature decrease in recent decade in both urban and rural stations may therefore be attributed to large background of

climate. The fact that the temperature decrease is more significant at Wuhan station than that at the rural stations leads to annual mean, minimum and maximum temperature UHII decrease during the nearly 10 years. The reasons for the more rapid decline of temperature at the urban station need to be further studied, but it is possible that the areas around the reference or suburban stations experience a more rapid urbanization than those around the urban station.

It is notable that the relative humidity (RH) has a certain recovery recently, and the RH series is inversely related to annual mean temperature series at Wuhan station, with the correlation coefficient reaching -0.65 for 1961–2015, and the urban heat island phenomenon is always accompanied by the urban dry island (Suonan et al.^[23]; Zhou^[43]). Rising of RH in the last 10 years therefore is not favorable for the formation or development of urban heat island around the urban station. Besides, sunshine hours have declined after 2007 in Wuhan^[23], and the winter mean wind speed near the ground surface has strengthened in the period of climate warming slowdown (Lin et al.^[44]). These, in combination with the background climate cooling of eastern China and the urbanization process around the rural stations, may also account for part of the greatly weakened UHII and the large downward trend of temperature during the last decade at the urban station.




Figure 7. Annual mean, minimum and maximum surface air temperature at Wuhan station and the two rural stations (average of two rural stations). Black solid line denotes 3 points moving average.

The inhomogeneity in the observed SAT series has a significant effect on the long-term trend analysis. In this paper, the breakpoints due to the station moving are detected for the urban and rural stations. There is a sudden drop in temperature after Wuhan station moved to the suburbs in 2010, with the T_{\min} difference being 1.35°C after the move. Historical data validates that this decrease is not a natural change, but an inhomogeneity error caused by the relocation (Liu et al.^[45]). It makes the T_{mean} trend drop from the $0.57^{\circ}\text{C}/10\text{yr}$ to $0.43^{\circ}\text{C}/10\text{yr}$. Meanwhile, the three kinds of temperature based UHII will be weaker after 2010, especially the mean and minimum temperature based UHII, if the inhomogeneity is not corrected. The effect of inhomogeneity caused by relocation on SAT series is obvious, and it is necessary to make reasonable adjustments.

Compared with previous studies on Wuhan station, this work tests and corrects the inhomogeneities of SAT data, and uses a more objective method to select rural stations. The SAT data series is also updated. The conclusions are very similar to those of previous works. The annual mean urbanization effect and contribution are $0.26^{\circ}\text{C}/10\text{yr}$ and 64.7% respectively in this analysis, like those estimated in Ren et al.^[13] ($0.20^{\circ}\text{C}/10\text{yr}$ and 64.5% for period 1961–2000), and Chen et al.^[20] ($0.24^{\circ}\text{C}/10\text{yr}$ and 60.4% for period 1960–2005). The previous studies also showed that urban warming is the highest in autumn and the second in summer, whereas summer witnesses the largest urbanization contribution, followed by autumn^[13], which is basically the same with the results of this paper. In addition, Wang et al. showed an annual mean urban warming of $0.31^{\circ}\text{C}/10\text{yr}$ for Wuhan station for the period 1961–2010, larger than the value of $0.19^{\circ}\text{C}/10\text{yr}$ for 1961–2015 in this paper^[34]. This difference may be mainly related to the updated data in this article, and the different rural stations selected in the two analyses.

The annual and seasonal mean urbanization effects have weakened when the data is updated to 2015, and their seasonal distribution has a slight change with the largest urbanization effect now occurring in summer and the second largest in autumn. This may be related to the increase in anthropogenic heat release due to the recent wider use of air conditioning and refrigeration equipment in summer (Zhou and Wen^[46]), as well as the compressed regulation effect of wetlands due to the rapid shrink of urban lake areas (Dan et al.^[47]).

4 CONCLUSIONS

The annual mean UHII is 0.39°C for 1961–2015 at Wuhan station, and the UHI effect is more obvious during the recent 28 years, with the annual mean UHII and minimum temperature based UHII reaching 0.69°C and 1.07°C respectively. The strongest monthly mean UHII appears in July, and the weakest in January.

The annual mean, minimum and maximum temperature based UHII shows significant upward trends for the period 1961–2015. The annual mean urban

warming and urbanization contribution are $0.18^{\circ}\text{C}/10\text{yr}$ and 48.8% respectively, and T_{\min} registers larger and more significant urban warming and urbanization contribution. The largest seasonal mean urban warming and urbanization contribution occurs in summer.

The annual mean urban warming becomes more obvious with time, with urbanization effect reaching as high as $0.53^{\circ}\text{C}/10\text{yr}$ for the period 1988–2004. The seasonal mean urbanization effect and urbanization contribution for the period 1988–2004 are both high in summer and autumn, with the greatest urban warming of higher than $1.00^{\circ}\text{C}/10\text{yr}$ seen for autumn mean T_{\min} .

From 2005 to 2015, the annual mean UHII for T_{mean} , T_{\min} and T_{max} show evident decreasing trends. This decrease may at least have been partially caused by the unrepresentativeness of the rural stations during the recent decade.

Therefore, the SAT in Wuhan station has been significantly affected by urbanization. The long-term observational data of surface air temperature at Wuhan station can be used to monitor and study urban climatology and urban climate change, but they are improper to be used in monitoring and studies of regional and global climate change without adjustment of the urbanization-induced bias, no matter whether the homogenization of the data is made or not.

REFERENCES:

- [1] JONES P D, GROISMAN P Y, COUGHLAN M, et al. Assessment of urbanization effects in time series of surface air temperature over land [J]. *Nature*, 1990, 347(6289): 171-177.
- [2] PETRESON T C. Assessment of urban versus rural in situ surface temperatures in the contiguous United States: No difference found [J]. *J Climate*, 2003, 16(18): 2941-2959.
- [3] PARKER D E. A demonstration that large-scale warming is not urban [J]. *J Climate*, 2006, 19(12): 2882-2895.
- [4] LAWRIMORE J H, MENNE M J, GLEASON B E, et al. An overview of the global historical climatology network monthly mean temperature data set, version 3 [J]. *J Geophys Res Atmos*, 2011, 116(D19): 121-139.
- [5] KARL T R, DIAZ H F, KUKLA G. Urbanization: Its detection and effect in the United States climate record [J]. *J Climate*, 1988, 1(11): 1099-1123.
- [6] HANSEN J, RUEDY R, SATO M, et al. A closer look at United States and global surface temperature change [J]. *J Geophys Res Atmos*, 2001, 106(D20): 23947-23963.
- [7] KALNAY E, CAI M. Impact of urbanization and land use on climate change [J]. *Nature*, 2003, -1(6939): 528-531.
- [8] ZHOU L, DICKINSON R E, TIAN Y, et al. Evidence for a significant urbanization effect on climate in China [C]. *Proceedings of the National Academy of Sciences of the United States of America*, 2004, 101(26): 9540-9544.
- [9] ZHOU Ya-qing, REN Guo-yu. Identifying and correcting urban bias for regional surface air temperature series of North China over period of 1961–2000 [J]. *Clim Environ Res*, 2005, 10(4): 743-753 (in Chinese).
- [10] REN G, ZHOU Y, CHU Z, et al. Urbanization effects on observed surface air temperature trends in north

- China [J]. *Jf Climate*, 2008, 21(6): 1333-1348.
- [11] XIE Zhuang, CUI Ji-liang, CHEN Da-gang, et al. The annual, seasonal and monthly characteristics of diurnal variation of urban heat island intensity in Beijing [J]. *Clim Environ Res*, 2006, 11(1): 69-75 (in Chinese).
- [12] ZHENG Zuo-fang, LIU Wei-dong, WANG Ying-chun, et al. Distributive character of urban heat island effect in the Beijing region [J]. *J Nanjing Inst Meteor*, 2006, 29 (5): 694-699 (in Chinese).
- [13] REN G, CHU Z, CHEN Z, et al. Implications of temporal change in urban heat island intensity observed at Beijing and Wuhan stations [J]. *Geophys Res Lett*, 2007, 34(5): 89-103.
- [14] YANG P, REN G, LIU W. Spatial and temporal characteristics of Beijing urban heat island intensity [J]. *J Appl Meteor Climatol*, 2013, 52(8): 1803-1816.
- [15] BIAN Tao, REN Guo-yu, ZHANG Cui-hua, et al. Characteristics and change of urban climate in Shijiazhuang City, Hebei Province, China [J]. *J Nanjing Uni Info Sci Technol: Nat Sci Ed*, 2012, 4 (5): 402-408 (in Chinese).
- [16] DENG Lian-tang, SHU Jiong, LI Chao-yi. Analysis on the changing characteristics of urban heat island in Shanghai [J]. *J Trop Meteor*, 2001, 17 (3): 273-280 (in Chinese).
- [17] LI Q, ZHANG H, LIU X, et al. Urban heat island effect on annual mean temperature during the last 50 years in China [J]. *Theor Appl Climatol*, 2004, 79(3-4): 165-174.
- [18] CHU Zi-ying, REN Guo-yu. Effect of enhanced urban heat island magnitude on average surface air temperature series in Beijing Region [J]. *J Meteor Res*, 2005, 63(4): 534-540 (in Chinese).
- [19] WU Yun, REN Yong-jian, FANG Si-da, et al. Analysis of the urban heat island effect in Wuhan and its influence elements [J]. *Meteor Disast Reduct Res*, 2017, 40(1): 43-50 (in Chinese).
- [20] CHEN Zheng-hong, WANG Hai-jun, REN Guo-yu. Study on asymmetric change of urban heat island intensity in Wuhan City [J]. *Adv Clim Change Res*, 2007, 3(5): 282-286 (in Chinese).
- [21] CAO Xiao-xue. Analysis on characteristics of climate change in Wuhan City form 1961-2012 [D]. Central China Normal University, 2015 (in Chinese).
- [22] FYFE J C, GILLET N P. Recent observed and simulated warming [J]. *Nature Clim Chang*, 2014, 4(3): 150-151.
- [23] SUONAN Kan-zhuo, REN Guo-yu, JIA Wen-qian, et al. Climatological characteristics and long-term trend of relative humidity in Wuhan City [J]. *Clim Environ Res*, 2018, 23(6): 715-724 (in Chinese).
- [24] GALLO K P, MCNAB A L, KARL T R, et al. The use of a vegetation index for assessment of the urban heat island effect [J]. *Int J Remote Sens*, 1993, 14 (11): 2223-2230.
- [25] REN Y, REN G. A remote-sensing method of selecting reference stations for evaluating urbanization effect on surface air temperature trends [J]. *J Climate*, 2011, 24(7): 3179-3189.
- [26] ZHANG Lei. Urbanization effect on extreme temperature change in East Asia [D]. Nanjing University of Information Science and Technology, 2014 (in Chinese).
- [27] WINKLER J A, SKAGGS R H, BAKER D G. Effect of temperature adjustments on the Minneapolis-St.Paul urban heat island [J]. *J Appl Meteor*, 1981, 20 (11): 1295-1300.
- [28] WANG Li-hong. Trend analysis of precipitation change in Wuhan City [J]. *Water Resour Power*, 2010, 28(12): 4-5 (in Chinese).
- [29] EASTERLING D R, PETERSON T C. A new method for detecting undocumented discontinuities in climatological time series [J]. *Int J Climatol*, 1995, 15(4): 369-377.
- [30] EASTERLING D R, PETERSON T C. The effect of artificial discontinuities on recent trends in minimum and maximum temperatures [J]. *Atmos Res*, 1995, 37 (1-3): 19-26.
- [31] LI Qing-xiang, LIU Xiao-ning, ZHANG Hong-zheng, et al. Homogeneity study of in situ observational climate series [J]. *Meteor Sci Technol*, 2003, 31 (1): 3-10 (in Chinese).
- [32] CAO Li-juan, YAN Zhong-wei. Research progress of homogeneity of ground climate data [J]. *Adv Clim Change Res*, 2011, 7(2): 129-135 (in Chinese).
- [33] WEI Feng-ying. Modern Climate Statistics Diagnosis and Prediction Technology [M]. Beijing: Meteorological Press, 2007 (in Chinese).
- [34] WANG Miao, WU Jun, REN Yong-jian, et al. An analysis of the features of the urban heat island in Wuhan and the influence elements [C]// 32th Annual Meeting of the Chinese Meteorological Society - S6 Addressing Climate Change, Low Carbon Development and Ecological Civilization Construction. China Meteorological Society, 2015: 14-28 (in Chinese).
- [35] ZHANG L, REN G, REN Y, et al. Effect of data homogenization on estimate of temperature trend: A case of Huairou Station in Beijing Municipality [J]. *Theor Appl Climatol*, 2014, 115(3-4): 365-373.
- [36] ZHOU Shu-zhen, ZHANG Chao. Heat island effect at Shanghai City [J]. *Acta Geograph Sinica*, 1982, 37(4): 372-382 (in Chinese).
- [37] XIAO Jian-ping, FENG Ping, LI Jian-yi. Maps testify the development track about the city of Wuhan during the decade [J]. *J Geomat*, 2014, 39(2): 59-62 (in Chinese).
- [38] BAI Hu-zhi, REN Guo-yu, FANG Feng. Characteristics of urban heat island effect and its influential factors in Lanzhou City [J]. *Meteor Sci Technol*, 2005, 33 (6): 492-495 (in Chinese).
- [39] CHEN Bing, SHI Guang-yu. Estimation of the distribution of global anthropogenic heat flux [J]. *Atmos Ocean Sci Lett*, 2012, 5(2): 108-112 (in Chinese).
- [40] XU Xiang-de. Introduction to urbanization and environmental meteorology [M]. Beijing: Meteorological Press, 2002 (in Chinese).
- [41] WANG Dai. Minimum temperature in China response to the warming hiatus [D]. Nanjing University of Information Science and Technology, 2017 (in Chinese).
- [42] XU Yan, TANG Guo-li, ZHANG Qiang. Analysis of the variation of the air temperature over China during the global warming hiatus period [J]. *Climate Change Reserch*, 2017, 13(6): 569-577 (in Chinese).
- [43] ZHOU Shu-zhen. Discussion of the urban-rural differences in atmospheric humidity of Shanghai area [J]. *Transact Oceanol Limnol*, 1994, (2): 13-25 (in Chinese).
- [44] LIN Chang-gui, YANG Kun, QIN Jun, et al. Observed coherent trends of surface and upper-air wind speed over China since 1960 [J]. *J Climate*, 2013, 26(9): 2891-2903.

- [45] LIU Wu-bi, LI Ming, JIANG Hong. Analysis of the impact of urban development on meteorological observation data in Wuhan [C]// The 33th Annual Meeting of the Chinese Meteorological Society - S10 Cities, Precipitation and Haze, China Meteorological Society, 2016: 6-12 (in Chinese).
- [46] ZHOU Jin, WEN Yuan-gao. Influence of UHI intensity on annual energy consumption for heating and air-conditioning in business buildings [J]. Building Tech Develop, 2012, 39(10): 58-59 (in Chinese).
- [47] DAN Yong-li, WANG Hong-zhi, ZHANG Huan, et al. Lakes evolution of central Wuhan during 2000 to 2010 [J]. Acta Ecolog Sinica, 2014, 34(5): 1311-1317 (in Chinese).

Citation: JIA Wen-qian, REN Guo-yu, SUONAN Kan-zhuo, et al. Urban heat island effect and its contribution to observed temperature increase at Wuhan station, central China [J]. J Trop Meteor, 2019, 25(1): 102-113.